arXiv:1302.3874v1 [hep-th] 15 Feb 2013

DAMTP-2013-9

Thermodynamics of “exotic” BTZ black holes

Paul K. Townsend[1,*] and Baocheng Zhang[2,†]

[1] *Department of Applied Mathematics and Theoretical Physics, Centre for Mathematical Sciences,*
University of Cambridge, Wilberforce Road, Cambridge, CB3 0WA, UK
[2] *State Key Laboratory of Magnetic Resonances and Atomic and Molecular Physics,*
Wuhan Institute of Physics and Mathematics, Chinese Academy of Sciences, Wuhan 430071, China

A number of three-dimensional (3D) gravity models, such as 3D conformal gravity, admit “exotic” black hole solutions: the metric is the same as the BTZ metric of 3D Einstein gravity but with reversed roles for mass and angular momentum, and an entropy proportional to the length of the inner horizon instead of the event horizon. Here we show that the BTZ solutions of the “exotic” 3D Einstein gravity (with parity-odd action but Einstein field equations) are exotic black holes, and we investigate their thermodynamics. The first and second laws of black hole thermodynamics still apply, and the entropy still has a statistical interpretation.

PACS numbers: 04.70.Dy, 04.60.Kz

In three spacetime dimensions (3D) the Einstein gravitational field equations imply that the metric has constant curvature and hence, in the case of a negative cosmological constant, that it is locally isometric to anti-de Sitter (adS) space. One solution is therefore the adS vacuum but there is also a two-parameter family of stationary black hole solutions, found by Banados, Teitelboim and Zanelli [1]; the BTZ metric takes the form

$$ds^2 = -N^2 dt^2 + N^{-2} dr^2 + r^2 \left(d\phi + N^\phi dt\right)^2 . \tag{1}$$

The functions N^2 and N^ϕ are

$$N^2 = -8Gm + \frac{r^2}{\ell^2} + \frac{16G^2 j^2}{r^2} , \qquad N^\phi = -\frac{4Gj}{r^2} , \tag{2}$$

where ℓ is the adS radius and G the 3D Newton constant. The parameters (m, j) can be interpreted as the mass M and angular momentum J of the black hole in units for which $\hbar = 1$; i.e. $M = m$ and $J = j$. The laws of black hole thermodynamics [2] apply, with the entropy given by the Bekenstein-Hawking formula [3, 4], which states (in the 3D context) that the entropy is 1/4 of the length of the event horizon in Planck units. There is even a statistical mechanics interpretation of this entropy [5, 6]; the microstates are those of a holographically dual conformal field theory (CFT) about which little is known except that its central charges are [7]

$$c_L = c_R = \frac{3\ell}{2G} . \tag{3}$$

The gravity description of the CFT is valid at large ℓ/G.

Because the BTZ metric is locally isometric to adS$_3$, it solves the field equations of *any* 3D gravity model admitting an adS$_3$ vacuum, and the mass and angular momentum may then be equal to some other linear combination of the parameters. Rather surprisingly, a number of 3D gravity models have been found [8–11] for which the roles of mass and angular momentum are *reversed*, in the sense that

$$M = j/\ell , \qquad J = \ell m . \tag{4}$$

Such BTZ black holes have been called “exotic”. The latest addition [12] to the class of 3D gravity models for which BTZ black holes are exotic is 3D conformal gravity [13, 14], for which the simplest action is the Lorentz-Chern-Simons (LCS) term, introduced in [15] in the context of “topologically massive gravity”. The central charges of the CFT holographically dual to generic 3D gravity models may be computed by a variety of methods [12, 16], and for models admitting exotic BTZ black holes it is found, e.g. for conformal 3D gravity, that

$$c_R = -c_L . \tag{5}$$

This is what one might expect of a gravity model with a parity-odd action.

Like the Kerr solution of the 4D Einstein equations, the BTZ metric has two Killing horizons, located at $r = r_\pm$, where $r_\pm$ are the zeros of N^2:

$$r_\pm = \sqrt{2G\ell\left(\ell m + j\right)} \pm \sqrt{2G\ell\left(\ell m - j\right)} . \tag{6}$$

We may assume without loss of generality that $j \geq 0$. We shall also assume that $\ell m \geq j$, to ensure the existence of an event horizon at $r = r_+$. Note that the parameters (m, j) can be simply expressed in terms of the radii of the inner and outer horizons:

$$\ell m = \frac{r_+^2 + r_-^2}{8G\ell} , \qquad j = \frac{r_+ r_-}{4G\ell} . \tag{7}$$

The entropy of BTZ black holes has also been computed by various means, which include holographic methods [17] and Wald’s Noether charge formula [18], or its extension to allow for parity-violation [19]. Applied to the original BTZ black hole, these methods give a result in agreement with the Bekenstein-Hawking formula, but applied to exotic BTZ black holes they give a different result: the entropy is proportional to the length $2\pi r_-$ of the *inner* horizon.

This result calls into question the validity of the laws of black hole thermodynamics. For example, a perturbation of a "normal" BTZ black hole will *decrease* the length of its inner horizon [20, 21]. This is not problematic in itself but it illustrates the point that Hawking's area theorem (length in 3D) need not apply to inner horizons, and as the entropy of an exotic BTZ black hole is proportional to the length of the inner horizon, it becomes unclear whether the second law of thermodynamics still applies. In an attempt to resolve some of these difficulties, Park proposed a new formula for the BTZ black hole entropy in the context of higher-derivative 3D gravity theories [22]. His proposal was supported by a computation of the statistical entropy but it conflicts with results obtained by other methods and implies various thermodynamic abnormalities, such as negative temperature.

Here we investigate these issues in the context of what must surely be the simplest 3D gravity model for which BTZ black holes are exotic. This is the *parity-odd* action for 3D Einstein gravity with negative cosmological constant that Witten called (coincidentally) "exotic" [23]. To contrast the normal and exotic versions of 3D Einstein gravity, we give here their respective Lagrangian 3-forms in terms of independent dreibein 1-forms e^a and Lorentz connection 1-forms ω^a ($a = 0, 1, 2$), and their torsion and curvature 2-form field strengths,

$$T^a = de^a + \epsilon^{abc}\omega_b e_c\,, \qquad R^a = d\omega^a + \frac{1}{2}\epsilon^{abc}\omega_b\omega_c\,, \quad (8)$$

where the exterior product of forms is implicit. In the normal case the Lagrangian 3-form is

$$L = \frac{1}{8\pi G}\left[e_a R^a - \frac{1}{6\ell^2}\epsilon^{abc}e_a e_b e_c\right]. \quad (9)$$

This is the well-known Einstein-Cartan formulation, so we do not elaborate further. The Lagrangian 3-form for the exotic theory is

$$L_E = \frac{\ell}{8\pi G}\left[\omega_a\left(d\omega^a + \frac{2}{3}\epsilon^{abc}\omega_b\omega_c\right) - \frac{1}{\ell^2}e_a T^a\right]. \quad (10)$$

The first term is the LCS term (except that the connection is an independent one). Although the action obtained by integration of L_E does not preserve parity, its parity transform is minus itself (i.e. it is parity-odd) so the field equations do preserve parity. Varying with respect to e^a, we get the torsion constraint $T^a = 0$, which allows us to solve for ω^a; using this in the equation obtained by varying ω^a, we recover the same Einstein equations that follow from the standard parity-even Lagrangian 3-form (9).

An explanation of how the exotic action was found will also explain why its BTZ black holes are exotic. The 3D adS isometry group is $SO(2,2) \cong Sl(2;\mathbb{R}) \times Sl(2;\mathbb{R})$. It was shown in [24] that the field equations of the $SO(2,2)$ Chern-Simons (CS) action are equivalent to the Einstein equations with negative cosmological constant. Actually, since one may consider any linear combination of the two CS terms for the $Sl(2;\mathbb{R})$ factors, for which the 1-form potentials are

$$A^a_\pm = e^a \pm \ell\omega^a\,, \quad (11)$$

the choice of $SO(2,2)$ Chern-Simons term is not unique, although there is a unique choice that yields a parity-preserving action. Because CS terms are intrinsically parity odd, and because the 1-forms A^a_+ and A^a_- are interchanged by parity, only the *difference* of the two CS terms preserves parity, and for a suitable overall constant this combination yields the standard action of 3D general relativity. This observation was first made by Achúcarro [25] and later by Witten [23], who also wrote down an explicit form for the parity-odd action obtained by taking the sum of the two $Sl(2;\mathbb{R})$ CS terms. Written in terms of the 1-forms (e^a, ω^a) this "exotic" version has the Lagrangian 3-form of (10).

Let us now examine the implications for BTZ black holes. For each of the two $Sl(2;\mathbb{R})$ subgroups of $SO(2,2)$, we can define a conserved charge $Q_\pm$ as the holonomy of an asymptotic $U(1)$ connection [26]. Using (11) one then finds, in the "normal" case, that

$$\ell M = Q_+ + Q_-\,, \qquad J = Q_+ - Q_-\,, \qquad \text{(normal)} \quad (12)$$

so parity leaves M unchanged but flips the sign of J, as expected. In contrast, the exotic case gives

$$\ell M = Q_+ - Q_-\,, \qquad J = Q_+ + Q_-\,, \qquad \text{(exotic)} \quad (13)$$

so the roles of ℓM and J are reversed. For the BTZ solution of "normal" 3D gravity we have $M = m$ and $J = j$, so we will find that $\ell M = j$ and $J = \ell m$ when we view it as a solution of the "exotic" theory. In other words, *BTZ black holes are "normal" as solutions of "normal' 3D Einstein gravity, and "exotic" as solutions of "exotic" 3D Einstein gravity.*

We now turn to black hole thermodynamics. The Hawking temperature T of the BTZ black hole and the angular velocity Ω of its event horizon are

$$T = \frac{r_+^2 - r_-^2}{2\pi r_+\ell^2}\,, \qquad \Omega = \frac{r_-}{\ell r_+}\,. \quad (14)$$

These expressions for the intensive thermodynamic variables are geometrical in the sense that they depend only on the location of the Killing horizons and are model-independent, i.e. independent of the particular field equations that are solved by the BTZ metric.

In contrast, the extensive thermodynamic variables are model-dependent. For generality, we shall consider the case in which the mass and angular momentum are given by

$$M = \alpha m + \gamma j/\ell\,, \qquad J = \alpha j + \gamma \ell m\,, \tag{15}$$

for constants (α, γ). The cases $(\alpha, \gamma) = (1, 0)$ and $(\alpha, \gamma) = (0, 1)$ correspond, respectively, to the normal and exotic BTZ black holes. In the "normal" case the second of the relations (7) tells us that the product of the lengths of the inner and outer horizons is independent of M, as expected on general grounds [27–30], but in the "exotic" case it is independent of J.

Let us now write the first law of black hole thermodynamics as

$$dM - \Omega dJ = T dS\,. \tag{16}$$

We claim that this law is satisfied for M and J given by (15) if, and only if, the entropy S is given by

$$S = \frac{\pi}{2G}\left(\alpha r_+ + \gamma r_-\right)\,. \tag{17}$$

In fact, given only M as a linear combination of m and j, the expressions for both J and S can be fixed by requiring the validity of the first law. This can be verified by computing dS in terms of dM and dJ using (6) and (14). When $(\alpha, \gamma) = (1, 0)$ the entropy reduces to the usual Bekenstein-Hawking formula, but when $(\alpha, \gamma) = (0, 1)$ it is given by the "exotic" formula $S = S_E \equiv \pi r_-/(2G)$; i.e. it is now 1/4 the length, in Planck units, of the *inner* horizon.

Next we show that the entropy, as given by (17), obeys the second law, at least for the quasi-stationary process of infalling matter. Let p be the 3-momentum of a particle falling through the event horizon. Since the event horizon is a Killing horizon for the Killing vector

$$\xi = \partial_t + \Omega \partial_\phi\,, \tag{18}$$

we have $-p \cdot \xi \geq 0$ at the horizon. This implies that the changes dM and dJ in the mass and angular momentum of the black hole satisfy the inequality $dM \geq \Omega dJ$, and hence that

$$\ell r_+ dM \geq r_- dJ\,. \tag{19}$$

Using the expressions (15) for the mass and angular momentum, and the formulas (6) for $r_\pm$, we find that

$$dS \geq 0\,. \tag{20}$$

This is true for any constants (α, γ), in particular it is true for the exotic case $(\alpha, \gamma) = (0, 1)$, even though the entropy is then proportional to the length of the inner horizon!

It should be noted that, in arriving at these results, we have used the angular momentum and temperature associated to the *event* horizon, not to the inner horizon. We are *not* discussing here the "thermodynamics of inner horizons".

Finally, we consider the statistical entropy of exotic BTZ black holes. In the high-temperature ($\beta \to 0$) limit, the partition function $Z(\beta) = \operatorname{tr} e^{-\beta H}$ of a CFT with Hamiltonian H can be approximated by the integral

$$Z(\beta) = \int_0^\infty d\Delta\, \rho(\Delta) e^{-\beta\Delta}\,, \tag{21}$$

where Δ is the eigenvalue of the Virasoro generator L_0, and $\rho(\Delta) = e^{S(\Delta)}$ is the (smoothed) density of states, equal to the exponential of the entropy function $S(\Delta)$. Using the fact that $S \propto \sqrt{\Delta}$, the partition function can be evaluated in a saddle point approximation. Then, using modular invariance, the result can be related to the low temperature ($\beta \to \infty$) limit, which is dominated by the ground state, with energy determined by the central charge. In this way Cardy found a large-Δ approximation to $\rho(\Delta)$, such that [31]

$$S(\Delta) \approx 2\pi\sqrt{c\Delta/6}\,, \tag{22}$$

where c is the central charge. Applying this formula to the left and right sectors of the Brown-Henneaux CFT with central charges $c_L = c_R = 3\ell/(2G)$, and using the relations

$$2\Delta_L = \ell M + J\,, \qquad 2\Delta_R = \ell M - J\,, \tag{23}$$

one finds that [5, 6]

$$S_L = \pi\sqrt{\frac{\ell\,(\ell M + J)}{2G}}\,, \qquad S_R = \pi\sqrt{\frac{\ell\,(\ell M - J)}{2G}}\,. \tag{24}$$

The sum $S = S_L + S_R$ equals the Bekenstein-Hawking entropy $\pi r_+/(2G)$.

In the exotic case we expect to have $c_R = -c_L$, as in the 3D conformal gravity case, because the action is parity odd. This means that we expect

$$c_L = \frac{3\ell}{2G}\,, \qquad c_R = -\frac{3\ell}{2G}\,. \tag{25}$$

We might also expect to have $2\Delta_L = \ell M_E + J_E$ and $2\Delta_R = J_E - \ell M_E$ because this is what one gets from (23) by setting $J = \ell M_E$ and $\ell M = J_E$. The weak cosmic censorship condition $J_E \geq \ell M_E$, needed for the existence of a horizon, then ensures that $\Delta_R > 0$. However, if this were true then we would have $c_R \Delta_R < 0$ and the Cardy formula would give an imaginary entropy.

Given that $c_R < 0$, the "exotic" version of the Brown-Henneaux CFT must be non-unitary, but this is only because we consider both the left and right sectors together. We may consider each sector in isolation because there is no interaction between them. Taken in isolation, there is no significance to the sign of the central charge c_R because we can change its sign from positive to negative by declaring that physical states have negative norm instead of positive norm, so that the energy is now bounded from above rather than from below. This is just a change of conventions; only the sign of c_R relative to c_L is convention-independent. We suggest that $c_R < 0$ for exotic 3D gravity theories precisely because the conventions for the right-movers are the reverse of the usual ones, in the sense just explained. This would flip the sign of Δ_R so that

$$2\Delta_L = \ell M_E + J_E \,, \qquad -2\Delta_R = J_E - \ell M_E \,. \tag{26}$$

Observe that the weak cosmic censorship condition $J_E - \ell M_E$ now ensures that $\Delta_R < 0$, and hence $c_R \Delta_R > 0$. Applying the Cardy formula (22) we find that

$$S_L = \pi\sqrt{\frac{\ell\,(J+\ell M)}{2G}}\,, \qquad S_R = \pi\sqrt{\frac{\ell\,(J-\ell M)}{2G}}\,. \tag{27}$$

Now we must ask how the partition function Z_R changes if we change the sign of the norm of all physical states so that $c_R \to -c_R$ and all energies change sign. We have seen that the laws of black hole thermodynamics continue to apply to exotic BTZ black holes, so we expect standard thermodynamic relations such as $E = -\partial \ln Z/\partial\beta$ to remain valid, but this relation is maintained when $E \to -E$ only if we also take $Z \to Z^{-1}$. We conclude from this that whereas $Z = Z_L Z_R$ in the "normal" case, we must have $Z = Z_L/Z_R$ in the exotic case, and this means that we should now *subtract* S_R from S_L to get the total "exotic" entropy:

$$S_E = \pi\sqrt{\frac{\ell\,(J+\ell M)}{2G}} - \pi\sqrt{\frac{\ell\,(J-\ell M)}{2G}} = \frac{\pi r_-}{2G}\,. \tag{28}$$

This is precisely the entropy required for the validity of the laws of black hole thermodynamics for exotic BTZ black holes.

In this paper, we have addressed some of the puzzles that have arisen in the study of the thermodynamics of "exotic" BTZ black holes, for which the three-dimensional spacetime is that of the usual BTZ black hole but with a reversal of the roles of mass and angular momentum. We have shown that such black holes occur even for 3D Einstein gravity in the context of its parity-odd "exotic" action. The relationship of the "normal" to the "exotic" versions of 3D Einstein gravity explains the reversal of the roles of mass and angular momentum. It also goes a long way to explaining why the BTZ black holes of conformal 3D gravity are exotic: it is because the exotic Einstein gravity can be viewed as a truncation of conformal 3D gravity; from the Chern-Simons perspective one is just restricting to an $SO(2,2)$ subgroup of $SO(2,3)$.

Our investigation of the thermodynamics of BTZ black holes allowed for the mass and angular momentum to be given by arbitrary linear combinations of the parameters of the BTZ metric. In fact, given only the mass, the first law determines the entropy, which satisfies the second law, at least for quasi-static perturbations. For exotic black holes this entropy is equal to 1/4 of the length of the *inner* horizon. Our derivation of this result, which agrees with earlier investigations, is general and applies to all models admitting exotic BTZ black holes. In fact, our entropy formula is valid for any BTZ black hole, so it also applies to higher-derivative 3D gravity models such as topologically massive gravity. A similar extension to higher-derivative 3D gravity of our arguments concerning the statistical entropy appears more problematic because the central charges of the dual CFT can change sign as the parameters of the model change. We leave this to future investigations.

Acknowledgements: PKT thanks Gary Gibbons and David Tong for helpful conversations. BZ is grateful for the hospitality of the General Relativity group in the Department of Applied Mathematics and Theoretical Physics of the University of Cambridge, where this work was begun, and he also acknowledges support from Grant No. 11104324 of the National Natural Science Foundation of China.

---

[*] Electronic address: p.k.townsend@damtp.cam.ac.uk
[†] Electronic address: zhangbc@wipm.ac.cn

[1] M. Bañados, C. Teitelboim and J. Zanelli, Phys. Rev. Lett. **69,** 1849 (1992).
[2] J. M. Bardeen, B. Carter and S. W. Hawking, Commun. Math. Phys. **31,** 161 (1973).
[3] J. D. Bekenstein, Phys. Rev. D **7,** 2333 (1973).
[4] S. W. Hawking, Commun. Math. Phys. **43,** 199 (1975); Erratum-ibid. **46,** 206 (1976).
[5] A. Strominger, JHEP **9802,** 009 (1998) [hep-th/9712251].
[6] D. Birmingham, I. Sachs and S. Sen, Phys. Lett. B **424** (1998) 275 [hep-th/9801019].
[7] J. D. Brown and M. Henneaux, Commun. Math. Phys. **104,** 207 (1986).
[8] S. Carlip and J. Gegenberg, Phys. Rev. D **44,** 424 (1991).
[9] S. Carlip, J. Gegenberg and R. B. Mann, Phys. Rev. D **51,** 6854 (1995).
[10] M. Bañados, Phys. Rev. D **57,** 1068 (1998).
[11] M. Bañados, A. Gomberoff and C. Martinez, Classical Quantum Gravity **15,** 3575 (1998).
[12] H. Afshar, B. Cvetkovic, S. Ertl, D. Grumiller and N. Johansson, Phys. Rev. D **85,** 064033 (2012).
[13] P. Van Nieuwenhuizen, "Three-dimensional Conformal Supergravity And Chern-simons Terms," In *Salam, A. (ed.), Sezgin, E. (ed.): Supergravities in diverse dimensions, vol. 2* 862-868
[14] J. H. Horne and E. Witten, Phys. Rev. Lett. **62,** 501 (1989).
[15] S. Deser, R. Jackiw and S. Templeton, Annals Phys. **140,** 372 (1982); Erratum-ibid. **185,** 406 (1988); Annals Phys. **185,** 406 (1988); Annals Phys. **281,** 409 (2000).

[16] P. Kraus and F. Larsen, JHEP **0601,** 022 (2006).

[17] S. N. Solodukhin, Phys. Rev. D **74,** 024015 (2006).

[18] R. M. Wald, Phys. Rev. D **48,** 3427 (1993).

[19] Y. Tachikawa, Classical Quantum Gravity **24,** 737 (2007).

[20] M. -I. Park, Classical Quantum Gravity **25,** 095013 (2008).

[21] S. Detournay, Phys. Rev. Lett. **109,** 031101 (2012).

[22] M. -I. Park, Phys. Lett. B **647,** 472 (2007); Phys. Rev. D **77,** 026011 (2008); Phys. Rev. D **77,** 126012 (2008).

[23] E. Witten, Nucl. Phys. B **311,** 46 (1988).

[24] A. Achúcarro and P. K. Townsend, Phys. Lett. B **180,** 89 (1986).

[25] A. Achúcarro, "Three-dimensional theories of Anti-de Sitter supergravity associated to the supergroups $\mathrm{OSp}(p/2;\mathrm{R}) \times \mathrm{OSp}(q/2;\mathrm{R})$", MSc Thesis, Univ. of the Basque Country, Bilbao, 1986.

[26] J. M. Izquierdo and P. K. Townsend, Classical Quantum Gravity **12,** 895 (1995).

[27] F. Larsen, Phys. Rev. D **56,** 1005 (1997).

[28] M. Cvetic, G. W. Gibbons and C. N. Pope, Phys. Rev. Lett. **106,** 121301 (2011).

[29] A. Castro and M. J. Rodriguez, Phys. Rev. D **86,** 024008 (2012).

[30] M. Ansorg and J. Hennig, Phys. Rev. Lett. **102,** 221102 (2009).

[31] J. L. Cardy, Nucl. Phys. B **270,** 186 (1986).